

Analysis and Discretization of Optimal Control Problems in Peridynamics

Joshua Siktar

jsiktar@vols.utk.edu

(joint with Tadele Mengesha and Abner J. Salgado)

2023 SIAM Central States Section (Theoretical and Applied Aspects of Nonlocal Models)

October 2023



THE UNIVERSITY OF
TENNESSEE
KNOXVILLE

DEPARTMENT OF
MATHEMATICS

Acknowledgments

- Thanks to my co-advisors Abner J. Salgado and Tadele Mengesha for their continued mentorship
- Thanks to NSF grant 2111228 for financial support
- Thanks to Animesh Biswas, Javier Cueto, and Scott Hootman Ng for organizing the session

- 1 Problem Formulation
- 2 Notation and Functional Setting
- 3 Well-posedness
- 4 Convergence of state equation as $\delta \rightarrow 0^+$
- 5 Discretization
- 6 Asymptotic compatibility

Problem statement

Find $(\overline{u}_\delta, \overline{g}_\delta) \in X_0 \times Z_{\text{ad}}$ such that

$$I(\overline{u}_\delta, \overline{g}_\delta) = \min_{g_\delta \in L^2, u_\delta \in X_0} \left\{ \int_{\Omega} F(x, u_\delta(x)) dx + \frac{\lambda}{2} \int_{\Omega} \Gamma(x) |g_\delta(x)|^2 dx \right\},$$

subject to the strongly coupled system of equations

$$\mathcal{L}_\delta u_\delta = g_\delta$$

- $\delta \geq 0$ is the degree of non-locality (the **horizon parameter** in peridynamics);
- \overline{g}_δ is a [vector-valued] external force;
- \overline{u}_δ represents the [vector-valued] displacement.
- \mathcal{L}_δ is a non-local operator, exact formula to be given.

Sample candidate integrand

Suppose we want to deform our material to fit a predetermined hole of shape u_{des} as closely as possible. Then our candidate integrand is

$$F(x, u(x)) = |u(x) - u_{\text{des}}(x)|^2$$



Material
(deformable)



Hole (fixed
shape)

Goals

- Show existence and uniqueness of minimizers
- Consider behavior as $\delta \rightarrow 0^+$
- Discretize via FEMs
- Study simultaneous limit as $\delta, h \rightarrow 0^+$ (asymptotic compatibility)

Bi-linear forms

Bi-linear form induces non-local state equation:

$$B_\delta(u, v) := \iint_{\mathcal{D}_\delta} \mathfrak{A}(x, y) k_\delta(x - y) \frac{Du(x, y)}{|x - y|} \frac{Dv(x, y)}{|x - y|} dx dy = \int_{\Omega} g(x) \cdot v(x) dx$$

- $\mathfrak{A}(x, y) = \frac{\alpha(x) + \alpha(y)}{2}$ is positive, bounded, symmetric
- k_δ a kernel with suitable properties
- Du is projected difference, $Du(x, y) := (u(x) - u(y)) \cdot \frac{x - y}{|x - y|}$
- $\Omega_\delta := \{x, \text{dist}(x, \partial\Omega) < \delta\} \cup \Omega$
- $\mathcal{D}_\delta = (\Omega \times \Omega_\delta) \cup (\Omega_\delta \times \Omega)$
- Models arise in peridynamics, specifically bond-based model

Function spaces

Our function space is based on B_δ :

$$X(\Omega_\delta; \mathbb{R}^n) := \{u|_\Omega \in L^2(\Omega; \mathbb{R}^n), B_\delta(u, u) < \infty\}$$

Version with zero non-local boundary data:

$$X_0(\Omega_\delta; \mathbb{R}^n) := \{u \in X(\Omega_\delta; \mathbb{R}^n), u = 0 \text{ in } \Omega_\delta \setminus \Omega\}$$

State equation is well-posed!

Proposition (Nonlocal Poincaré-Korn (Mengesha-Du 2014))

There exists a $\delta_0 > 0$ and a constant $C(\delta_0) > 0$ such that for all $\delta \in (0, \delta_0]$ and $u \in X_0(\Omega_\delta; \mathbb{R}^n)$,

$$\|u\|_{L^2(\Omega; \mathbb{R}^n)}^2 \leq C(\delta_0) \int_{\Omega_\delta} \int_{\Omega_\delta} k_\delta(x-y) \frac{|Du(x,y)|^2}{|x-y|^2} dx dy.$$

NOTE: The projected difference vanishes under infinitesimal rigid displacements.

Theorem (Existence and Uniqueness for State Equation)

For any $g_\delta \in L^2$, there exists a unique $u_\delta \in X_0$ such that the state system

$$B_\delta(u_\delta, w_\delta) = \langle g_\delta, w_\delta \rangle$$

is satisfied for all $w_\delta \in X_0$. Furthermore, we have the stability estimate

$$\|u_\delta\|_{X(\Omega_\delta; \mathbb{R}^n)} \lesssim \|g_\delta\|_{L^2(\Omega_\delta; \mathbb{R}^n)}$$

for some constant independent of δ .

Cost functional assumptions

$$I(u, g) := \int_{\Omega} F(x, u(x)) dx + \frac{\lambda}{2} \int_{\Omega} \Gamma(x) |g(x)|^2 dx$$

Here Z_{ad} is a box constraint, and $F : \Omega \times \mathbb{R}^n \rightarrow \mathbb{R}$ is such that:

- 1 For all $v \in \mathbb{R}^n$ the mapping $x \mapsto F(x, v)$ is measurable;
- 2 For all $x \in \Omega$ the mapping $v \mapsto F(x, v)$ is continuous and convex;
- 3 There exist $c_1 > 0$ and $\ell \in L^1(\Omega)$

$$|F(x, v)| \leq c_1 |v|^2 + \ell(x)$$

for all $x \in \Omega$, $v \in \mathbb{R}^n$. Moreover, $\lambda \geq 0$, and $\Gamma \in L^\infty(\Omega)$ is positive a.e.

Well-posedness of optimal control problem

Theorem (Well-posedness)

For any $\delta > 0$: there exists $(\bar{u}_\delta, \bar{g}_\delta) \in X_0(\Omega_\delta; \mathbb{R}^n) \times Z_{ad}$ minimizing

$$I(u_\delta, g_\delta) = \int_{\Omega} F(x, u_\delta(x)) dx + \frac{\lambda}{2} \int_{\Omega} \Gamma(x) |g_\delta(x)|^2 dx,$$

where $\bar{u}_\delta \in X_0$ solves

$$B_\delta(u_\delta, v_\delta) = \int_{\Omega} g_\delta(x) \cdot v_\delta(x) dx \quad \forall v_\delta \in X_0$$

Furthermore, if F is strictly convex or $\lambda > 0$, then the minimizer is unique.

Convergence of state equation as $\delta \rightarrow 0^+$

Local bi-linear form (of Navier-Lamé system of linear elasticity)

$$B_0(u, v) := C(n) \int_{\Omega} \alpha(x) (2\langle \text{Sym}(\nabla u), \text{Sym}(\nabla v) \rangle_F + \text{div}(u)\text{div}(v)) dx$$

where $C(n) := \frac{1}{(n+2)(n+4)}$.

Theorem

Suppose $\{(\bar{u}_\delta, \bar{g}_\delta)\}_{\delta>0}$ is the family of solutions to the non-local problem. Then, there is (\bar{u}, \bar{g}) such that $\bar{u}_\delta \rightarrow \bar{u}$ in $L^2(\Omega; \mathbb{R}^n)$ and $\bar{g}_\delta \rightarrow \bar{g}$ in $L^2(\Omega; \mathbb{R}^n)$. Moreover, (\bar{u}, \bar{g}) solves the local optimal control problem.

Local state equation that (\bar{u}, \bar{g}) satisfies the constraint:

$$B_0(u, v) = \int_{\Omega} g(x) \cdot v(x) dx$$

Discrete Formulation

- Use continuous piecewise linear for states
- Use piecewise constants for controls
- Discrete optimal control problem is well-posed

Optimality conditions

Let S_δ denote solution operator for state equation

Non-local continuous optimality conditions (including adjoint):

$$\begin{aligned}\bar{u}_\delta &= S_\delta \bar{g}_\delta \\ \bar{p}_\delta &= S_\delta^* F_u(\cdot, \bar{u}_\delta) = S_\delta F_u(\cdot, \bar{u}_\delta) \\ \langle \bar{p}_\delta + \lambda \bar{g}_\delta, \gamma_z - \bar{g}_\delta \rangle_{L^2} &\geq 0, \quad \forall \gamma_z \in Z_{\text{ad}}.\end{aligned}$$

Projection formulas:

$$\begin{aligned}\bar{g}_\delta(x) &= \mathbb{P}_{Z_{\text{ad}}} \left(-\frac{1}{\lambda} \bar{p}_\delta(x) \right) \\ \bar{g}_{\delta,h}(x) &= \mathbb{P}_{Z_{\text{ad}}} \left(-\frac{1}{\lambda} \Pi_0 \bar{p}_{\delta,h}(x) \right)\end{aligned}$$

Here Π_0 is defined as $\Pi_{0T} z := f_T z$ on each triangle T in our mesh

Non-local problem convergence: summary

Theorem (Convergence)

In the setting of our problems,

$$\|\overline{u_\delta} - \overline{u_{\delta,h}}\|_{X(\Omega_\delta; \mathbb{R}^n)} \lesssim \omega(h) + \inf_{v_{\delta,h} \in X_{\delta,h}} \|\widehat{u}_\delta - v_{\delta,h}\|_{X(\Omega_\delta; \mathbb{R}^n)} + \inf_{v_{\delta,h} \in X_{\delta,h}} [\overline{u_\delta} - v_{\delta,h}]_{X(\Omega_\delta; \mathbb{R}^n)} + \inf_{v_{\delta,h} \in X_{\delta,h}} [\overline{p_\delta} - v_{\delta,h}]_{X(\Omega_\delta; \mathbb{R}^n)};$$

$$\|\overline{g_\delta} - \overline{g_{\delta,h}}\|_{L^2(\Omega; \mathbb{R}^n)}^2 \lesssim \omega(h)^2 + \left(\inf_{v_{\delta,h} \in X_{\delta,h}} [\overline{u_\delta} - v_{\delta,h}]_{X(\Omega_\delta; \mathbb{R}^n)} \right)^2 + \left(\inf_{v_{\delta,h} \in X_{\delta,h}} [\overline{p_\delta} - v_{\delta,h}]_{X(\Omega_\delta; \mathbb{R}^n)} \right)^2.$$

$$\|\overline{p_\delta} - \overline{p_{\delta,h}}\|_{X(\Omega_\delta; \mathbb{R}^n)} \lesssim \omega(h) + \inf_{v_{\delta,h} \in X_{\delta,h}} \|\widehat{p}_h - v_{\delta,h}\|_{X(\Omega_\delta; \mathbb{R}^n)} + \inf_{v_{\delta,h} \in X_{\delta,h}} \|\widehat{u}_\delta - v_{\delta,h}\|_{X(\Omega_\delta; \mathbb{R}^n)} + \inf_{v_{\delta,h} \in X_{\delta,h}} [\overline{u_\delta} - v_{\delta,h}]_{X(\Omega_\delta; \mathbb{R}^n)} + \inf_{v_{\delta,h} \in X_{\delta,h}} [\overline{p_\delta} - v_{\delta,h}]_{X(\Omega_\delta; \mathbb{R}^n)}.$$

Non-local problem convergence: comments

- $X_{\delta,h}$ is our discrete non-local space
- Here $\omega(h)$ represents the approximation properties of the projection Π_0
- If our kernel is of fractional type, i.e. $k_\delta(\xi) \sim \frac{1}{|\xi|^{n+2s}}$, then $\omega(h) \sim h^s$

Discrete analogue of convergence

Theorem (Discrete Convergence)

Suppose $\{(\overline{u}_{\delta,h}, \overline{g}_{\delta,h})\}_{\delta>0}$ is the family of solutions to the non-local discrete problem. Then, there is $(\overline{u}_h, \overline{g}_h)$ such that $\overline{u}_{\delta,h} \rightarrow \overline{u}_h$ in $L^2(\Omega; \mathbb{R}^n)$ and $\overline{g}_{\delta,h} \rightarrow \overline{g}_h$ in $L^2(\Omega; \mathbb{R}^n)$. Moreover, $(\overline{u}_h, \overline{g}_h)$ solves the local discrete optimal control problem.

What is asymptotic compatibility?


- Introduced by X. Tian and Q. Du (2014)
- Originally developed for linear, non-local state equations

$$\mathcal{L}_{\delta,h}u_{\delta,h} = f$$

- Unconditional convergence of approximations in both discretization and horizon parameters

Definition (Asymptotic Compatibility)

Given fixed data f in a Hilbert Space, the family of solutions $\{u_{\delta,h}\}_{\delta,h>0}$ is **asymptotically compatible** in $\delta, h > 0$ if for any sequences $\{\delta_k\}_{k=1}^{\infty}, \{h_k\}_{k=1}^{\infty}$ with $\delta_k, h_k \rightarrow 0$, we have that $u_{\delta_k, h_k} \rightarrow u_0$ strongly in some Hilbert space norm, where u_0 is the solution to a local, continuous problem.



What is asymptotic compatibility? (continued)

Definition (Asymptotic Compatibility for Optimal Control)

We say that the family of solutions $\{(\overline{u}_{\delta,h}, \overline{g}_{\delta,h})\}_{h>0, \delta>0}$ to the nonlocal discrete optimal control problem is **asymptotically compatible** in $\delta, h > 0$ if for any sequences $\{\delta_k\}_{k=1}^{\infty}, \{h_k\}_{k=1}^{\infty}$ with $\delta_k, h_k \rightarrow 0$, we have that $\overline{g}_{\delta_k, h_k} \rightharpoonup \overline{g}$ weakly in $L^2(\Omega; \mathbb{R}^n)$, and $\overline{u}_{\delta_k, h_k} \rightarrow \overline{u}$ strongly in $L^2(\Omega; \mathbb{R}^n)$. Here $(\overline{u}, \overline{g}) \in H_0^1(\Omega; \mathbb{R}^n) \times Z_{\text{ad}}$ denotes the optimal solution for the local continuous problem.

NOTE: In particular, $H_0^1(\Omega; \mathbb{R}^n)$ is the limiting space of $X_0(\Omega_\delta; \mathbb{R}^n)$ as $\delta \rightarrow 0^+$

$$\begin{array}{ccc}
 (\overline{u}_{\delta,h}, \overline{g}_{\delta,h}) & \xrightarrow{\delta \rightarrow 0^+} & (\overline{u}_h, \overline{g}_h) \\
 \downarrow h \rightarrow 0^+ & \searrow k \rightarrow \infty & \downarrow h \rightarrow 0^+ \\
 (\overline{u}_\delta, \overline{g}_\delta) & \xrightarrow{\delta \rightarrow 0^+} & (\overline{u}, \overline{g})
 \end{array}$$

Asymptotic compatibility: result and notation

Theorem

Assume \mathfrak{A} is Lipschitz. Then, our family of optimal control problems is asymptotically compatible as $\delta, h \rightarrow 0^+$.

Define $A_\delta : X_0(\Omega_\delta; \mathbb{R}^n) \rightarrow (X_0(\Omega_\delta; \mathbb{R}^n))^*$ as the invertible operator such that

$$\langle A_\delta u, v \rangle_{X_0^*, X_0} = B_\delta(u, v) \quad \forall u, v \in X_0(\Omega_\delta; \mathbb{R}^n);$$

define $A_0 : H_0^1(\Omega; \mathbb{R}^n) \rightarrow H^{-1}(\Omega; \mathbb{R}^n)$ as the invertible operator such that

$$\langle A_0 u, v \rangle_{H^{-1}, H_0^1} = B_0(u, v) \quad \forall u, v \in H_0^1(\Omega; \mathbb{R}^n).$$

Proof of asymptotic compatibility

Needed preliminaries:

- 1 Given a $v \in H_0^1(\Omega; \mathbb{R}^n)$, and $h_k, \delta_k \rightarrow 0$, we can find a sequence $v_k \in X_{\delta_k, h_k}$ such that $v_k \rightarrow v$ strongly in $H^1(\Omega; \mathbb{R}^n)$ as $k \rightarrow \infty$.
- 2 For any sequences $\{\delta_k\}_{k=1}^\infty, \{h_k\}_{k=1}^\infty$ with $\delta_k, h_k \rightarrow 0$, there exists a $C > 0$ so that $\|\overline{u_{\delta_k, h_k}}\|_{X(\Omega_{\delta_k}; \mathbb{R}^n)} \leq C$ uniformly in $k \in \mathbb{N}^+$.
- 3 We have that $A_\delta u \in L^2(\Omega; \mathbb{R}^n)$ and $\lim_{\delta \rightarrow 0^+} \|A_\delta u - A_0 u\|_{L^2(\Omega; \mathbb{R}^n)} = 0$ for each $u \in C_0^\infty(\Omega; \mathbb{R}^n)$.

Pick sub-sequence of $\{(\overline{u_k}, \overline{g_k}, \overline{p_k})\}_{k=1}^\infty$ so there is a limit point $(u_*, g_*, p_*) \in H_0^1(\Omega; \mathbb{R}^n) \times Z_{\text{ad}} \times H_0^1(\Omega; \mathbb{R}^n)$ with convergence in appropriate topology

Proof of asymptotic compatibility (continued)

Step 1: Show that $B_0(u_*, \varphi) = \langle g_*, \varphi \rangle$ for all $\varphi \in H_0^1(\Omega; \mathbb{R}^n)$

Pick $\varphi \in C_0^\infty(\Omega; \mathbb{R}^n)$, let $w_k := I_k \varphi$ (nodal interpolation), $w_k \rightarrow \varphi$ in $W^{1,\infty}(\Omega; \mathbb{R}^n)$, then compute limit as $k \rightarrow \infty$ of

$$B_{\delta_k}(\overline{u_k}, w_k) = \langle A_{\delta_k} \varphi, \overline{u_k} \rangle_{X_0^*, X_0} + \langle A_{\delta_k}(w_k - \varphi), \overline{u_k} \rangle_{X_0^*, X_0} =: I_k + II_k.$$

Step 2: Show that $B_0(\varphi, p_*) = \langle u_*, \varphi \rangle$ for all $\varphi \in H_0^1(\Omega; \mathbb{R}^n)$

Strategy is identical to Step 1!

Step 3: Show that $g_*(x) = -\frac{1}{\lambda} \mathbb{P}_{Z_{\text{ad}}}(p_*(x))$

Recall that $\overline{g_k}(x) = -\frac{1}{\lambda} \mathbb{P}_{Z_{\text{ad}}}(\Pi_0 \overline{p_k}(x))$, show $\Pi_0 \overline{p_k} \rightarrow p_*$ strongly in $L^2(\Omega; \mathbb{R}^n)$

Proof of asymptotic compatibility (continued)

Step 4: Unraveling

- Steps 1-3 and uniqueness of solutions to optimality system give $\bar{u} = u_*$, $\bar{g} = g_*$, and $\bar{p} = p_*$
- This is the limit point reached for any sub-sequence of original sequence
- Entire sequence of triples $\{(\bar{u}_k, \bar{g}_k, \bar{p}_k)\}_{k=1}^{\infty}$ converges to $(\bar{u}, \bar{g}, \bar{p})$

Recap and closing

- Showed existence and uniqueness of minimizers
- Considered behavior as $\delta \rightarrow 0^+$
- Discretized via FEMs
- Studied simultaneous limit as $\delta, h \rightarrow 0^+$

Thank you! Questions?

Auxiliary lemmas

Lemma (Regularity of Control for Fractional-Type Kernels)

Suppose that

$$\frac{c}{|\xi|^{n+2s}} \leq \frac{k_\delta(\xi)}{|\xi|^2} \leq \frac{C}{|\xi|^{n+2s}}$$

holds for all $\xi \in B(0, \delta)$, for some $s \neq \frac{1}{2}$. Then necessarily $\overline{g}_\delta \in X(\Omega_\delta; \mathbb{R}^n)$.

Recall Fractional Sobolev Space (special case of our kernels):

$$H^s(\Omega_\delta; \mathbb{R}^n) := \left\{ u|_\Omega \in L^2(\Omega; \mathbb{R}^n), \frac{|u(x) - u(y)|}{|x - y|^{\frac{n}{2} + s}} \in L^2(\Omega_\delta \times \Omega_\delta) \right\}$$

Proof strategy:

- Mengesha-Du 2016 says that $H^s(\Omega_\delta; \mathbb{R}^n) = X(\Omega_\delta; \mathbb{R}^n)$ here
- Use projection formula $\overline{g}_\delta(x) = -\frac{1}{\lambda} \mathbb{P}_{Z_{\text{ad}}}(\overline{p}_\delta(x))$
- This pointwise projection is continuous in H^s semi-norm but not X semi-norm!

Auxiliary lemmas (continued)

Lemma (Approximation)

If $w \in L^2(\Omega_\delta; \mathbb{R}^n)$, then

$$\|\Pi_0 w - w\|_{L^2(\Omega; \mathbb{R}^n)} \leq \omega(h),$$

where $\Pi_0 : L^2(\Omega_\delta; \mathbb{R}^n) \rightarrow Z_h$ denotes the projection of a function onto the piecewise constants with respect to the given mesh. If in fact $w \in X(\Omega_\delta; \mathbb{R}^n)$ and k_δ satisfies the fractional inequality on $B(0, \delta)$ for some $s \neq \frac{1}{2}$, then

$$\|\Pi_0 w - w\|_{L^2(\Omega; \mathbb{R}^n)} \lesssim h^s \|w\|_{X(\Omega_\delta; \mathbb{R}^n)}.$$

Control convergence (continued)

Galerkin Approximations:

$q_{\delta,h} \in X_{\delta,h}$ be the Galerkin approximation to \bar{p}_δ , i.e., the solution of

$$B_\delta(v_{\delta,h}, q_{\delta,h}) = \langle \bar{u}_\delta, v_{\delta,h} \rangle \quad \forall v_{\delta,h} \in X_{\delta,h}.$$

$U_{\delta,h} \in X_{\delta,h}$ for \bar{u}_δ :

$$B_\delta(U_{\delta,h}, v_{\delta,h}) = \langle \bar{g}_\delta, v_{\delta,h} \rangle \quad \forall v_{\delta,h} \in X_{\delta,h}.$$

$r_{\delta,h} \in X_{\delta,h}$ solves

$$B_\delta(v_{\delta,h}, r_{\delta,h}) = \langle U_{\delta,h}, v_{\delta,h} \rangle \quad \forall v_{\delta,h} \in X_{\delta,h}.$$

 Back

Control convergence (continued)

Let $l_1 := \langle \overline{p}_\delta - \overline{p}_{\delta,h}, \overline{g}_{\delta,h} - \overline{g}_\delta \rangle$ and $l_2 := \langle \overline{p}_{\delta,h} + \lambda \overline{g}_{\delta,h}, \Pi_0 \overline{g}_\delta - \overline{g}_\delta \rangle$. Using optimality conditions gives

$$\lambda \|\overline{g}_\delta - \overline{g}_{\delta,h}\|_{L^2(\Omega; \mathbb{R}^n)}^2 \leq l_1 + l_2.$$

$$l_1 = \langle \overline{p}_\delta - q_{\delta,h}, \overline{g}_{\delta,h} - \overline{g}_\delta \rangle + \langle q_{\delta,h} - r_{\delta,h}, \overline{g}_{\delta,h} - \overline{g}_\delta \rangle + \langle r_{\delta,h} - \overline{p}_{\delta,h}, \overline{g}_{\delta,h} - \overline{g}_\delta \rangle =: l_{1,1} + l_{1,2} + l_{1,3}.$$

By use of Galerkin approximations, find that $l_{1,3} \leq 0$ and

$$l_{1,1} \lesssim \|\overline{g}_{\delta,h} - \overline{g}_\delta\|_{L^2(\Omega; \mathbb{R}^n)} \inf_{v_{\delta,h} \in X_{\delta,h}} [\overline{p}_\delta - v_{\delta,h}]_{X(\Omega_\delta; \mathbb{R}^n)};$$

$$l_{1,2} \lesssim \|\overline{g}_{\delta,h} - \overline{g}_\delta\|_{L^2(\Omega; \mathbb{R}^n)} \inf_{v_{\delta,h} \in X_{\delta,h}} [\overline{u}_\delta - v_{\delta,h}]_{X(\Omega_\delta; \mathbb{R}^n)}.$$

Control convergence (continued)

By Young's Inequality,

$$I_1 \leq \frac{\lambda}{3} \|\overline{g}_{\delta,h} - \overline{g}_{\delta}\|_{L^2(\Omega; \mathbb{R}^n)}^2 + C \left(\inf_{v_{\delta,h} \in X_{\delta,h}} [\overline{u}_{\delta} - v_{\delta,h}]_{X(\Omega_{\delta}; \mathbb{R}^n)} \right)^2 + C \left(\inf_{v_{\delta,h} \in X_{\delta,h}} [\overline{p}_{\delta} - v_{\delta,h}]_{X(\Omega_{\delta}; \mathbb{R}^n)} \right)^2.$$

Control convergence (continued)

Now we write l_2 as

$$\begin{aligned} \langle \overline{p_{\delta,h}} + \lambda \overline{g_{\delta,h}}, \Pi_0 \overline{g_{\delta}} - \overline{g_{\delta}} \rangle &= \langle \overline{p_{\delta}} + \lambda \overline{g_{\delta}}, \Pi_0 \overline{g_{\delta}} - \overline{g_{\delta}} \rangle + \lambda \langle \overline{g_{\delta,h}} - \overline{g_{\delta}}, \Pi_0 \overline{g_{\delta}} - \overline{g_{\delta}} \rangle + \\ &\langle \overline{p_{\delta,h}} - r_{\delta,h}, \Pi_0 \overline{g_{\delta}} - \overline{g_{\delta}} \rangle + \langle r_{\delta,h} - q_{\delta,h}, \Pi_0 \overline{g_{\delta}} - \overline{g_{\delta}} \rangle + \langle q_{\delta,h} - \overline{p_{\delta}}, \Pi_0 \overline{g_{\delta}} - \overline{g_{\delta}} \rangle =: \\ &l_{2,1} + l_{2,2} + l_{2,3} + l_{2,4} + l_{2,5}. \end{aligned}$$

Control convergence (continued)

Use that $\Pi_0(\bar{p}_\delta + \lambda \bar{g}_\delta) = 0$ to estimate $l_{2,1}$ as

$$l_{2,1} \leq \omega(h).$$

For $l_{2,2}$, use Cauchy and stability:

$$l_{2,2} \leq \frac{\lambda}{3} \|\bar{g}_{\delta,h} - \bar{g}_\delta\|_{L^2(\Omega; \mathbb{R}^n)}^2 + \omega(h),$$

Control convergence (continued)

For $l_{2,3}$, use Galerkin approximations and the stability:

$$l_{2,3} \leq \omega(h)^2 + \left(\inf_{v_{\delta,h} \in X_{\delta,h}} [\bar{u}_{\delta} - v_{\delta,h}]_{X(\Omega_{\delta}; \mathbb{R}^n)} \right)^2$$

For $l_{2,4}$ and $l_{2,5}$, use Céa's lemma and Cauchy:

$$l_{2,4} \leq \omega(h)^2 + \left(\inf_{v_{\delta,h} \in X_{\delta,h}} [\bar{u}_{\delta} - v_{\delta,h}]_{X(\Omega_{\delta}; \mathbb{R}^n)} \right)^2 ;$$

$$l_{2,5} \leq \omega(h)^2 + \left(\inf_{v_{\delta,h} \in X_{\delta,h}} [\bar{p}_{\delta} - v_{\delta,h}]_{X(\Omega_{\delta}; \mathbb{R}^n)} \right)^2 .$$

Control convergence (continued)

Use Young's Inequality and combine all the estimates to get

$$\frac{\lambda}{3} \|\overline{g_\delta} - \overline{g_{\delta,h}}\|_{L^2(\Omega; \mathbb{R}^n)}^2 \lesssim \omega(h)^2 + \left(\inf_{v_{\delta,h} \in X_{\delta,h}} [\overline{u_\delta} - v_{\delta,h}]_{X(\Omega_\delta; \mathbb{R}^n)} \right)^2 + \left(\inf_{v_{\delta,h} \in X_{\delta,h}} [\overline{p_\delta} - v_{\delta,h}]_{X(\Omega_\delta; \mathbb{R}^n)} \right)^2.$$

 Back